


RESEARCH REPORT

RR-012



DYNAMICAL PARAMETERS DERIVED FROM ANALYTICAL FUNCTIONS  
REPRESENTING INDIAN MONSOON FLOW

by

S.T. AWADE & G.C. ASNANI

INDIAN INSTITUTE OF TROPICAL METEOROLOGY

Ramdurg House  
Ganeshkhind Road,  
Poona-5, India

November 1973

Dynamical parameters derived from analytical functions represent-  
ing Indian Monsoon Flow

by

S.T.Awade & G.C.Asnani

RB-12

Indian Institute of Tropical Meteorology, Poona-5, India.

Abstract

Normals of July upper wind (zonal component only) along longitude  $77\frac{1}{2}^{\circ}$  E were considered between latitudes  $5^{\circ}$  N and  $25^{\circ}$  N, at isobaric levels from 850 mb to 150 mb <sup>at</sup> interval of 50 mb. Parabola of the type  $u = a + b \log_{10} P + c (\log_{10} P)^2$  was fitted by the least square method to the zonal wind at each latitude at an interval of  $2\frac{1}{2}$  degree latitude. To the values of a, b and c at each latitude, a parabola of the type  $a = a_0 + a_1 \phi + a_2 \phi^2$  was fitted. Here u is zonal wind in kts, p is pressure in mb and  $\phi$  is latitude in degrees. The fitted values of u agree well with observed values.

Assuming geostrophic approximation through thermal wind relationship, meridional profile of temperature at each isobaric level has been determined in analytical form and compared with observed temperatures. Assumption of geostrophic balance in respect of climatological zonal winds appears to be justified. The method therefore gives detailed structure of temperature field and other parameters in a quantitative form which is otherwise difficult to build through ordinary analysis of temperature values at individual stations.

Static stability, vertical shear and absolute vorticity on isobaric and on isentropic surfaces are calculated with the help of these values. It is found that the zonal current is inertially stable and also



barotropically stable along  $77\frac{1}{2}^{\circ}$  E between  $5^{\circ}$  N and  $25^{\circ}$  N.

The analysis of maximum wind suggests that the easterly jet maximum is perhaps close to latitude  $9^{\circ}$  N.

## 1. Introduction

Wind shears in the vertical and in the horizontal, temperature gradient in meridional direction, static stability, Richardson number and absolute vorticity are some of the parameters essential for understanding of the dynamics of zonal current (Charney 1947, Fjortoft 1950, Kuo 1956 etc.). It is possible to calculate these parameters at individual stations straightaway from observations, but to get smooth values representing large scale features, it is better to have recourse to smoothing process. We feel that to get smooth profile, it is better to fit an analytical function to one basic parameter and to derive all other parameters from this function. We choose zonal wind as a basic parameter and derive all other quantities from the distribution of this zonal wind.

July is a representative monsoon month for the Indian region, The region of India and neighbourhood from equator to  $25^{\circ}$  N is fairly free from high mountain and is also well in the grip of summer monsoon. We consider this month suitable for the study of the zonal current. Longitude  $77\frac{1}{2}^{\circ}$  E was taken as representative of the Indian region.

We have taken the layer from 850 mb to 150 mb for the purpose of fitting a curve in the vertical. Easterly jet stream maximum occurs between 150 mb and 100 mb. We thought that inclusion of 150 to 100 mb layer along with that from 850 to 150 mb layer will necessitate the,



fitting of a very complicated curve. Aiming at simplicity of a curve, we confined ourselves to the layer from 850 to 150 mb.

From the analysis of the flow pattern, we found that the latitudinal belt from equator to  $5^{\circ}\text{N}$  could not be easily taken along with the region north of  $5^{\circ}\text{N}$ . While we fitted analytical function north of  $5^{\circ}\text{N}$ , we used only graphical interpolation from  $5^{\circ}\text{N}$  to equator.

However, in fixing the value of zonal wind from  $5^{\circ}\text{N}$  to equator, we were some what guided by the consideration that at the levels of weak meridional flow, the relative vorticity of zonal wind may be kept as zero at the equator and absolute vorticity may not become negative north of equator.

Thus, we have fitted analytical functions from  $5^{\circ}\text{N}$  to  $25^{\circ}\text{N}$  and from 850 mb to 150 mb, along the longitude  $77\frac{1}{2}^{\circ}\text{E}$ , during the month of July. We adopted the graphical method of smoothing from  $5^{\circ}\text{N}$  to equator.

## 2. Data and computational procedure

For the present study, "Normals (1965) of Rawin winds based on afternoon data", issued by the office of the DDGC, India Met. Department, were the main source of data.

### 2.1 Fitting a function to the zonal wind

The zonal component  $u$  of the normal wind was plotted at each isobaric level for the Indian region. Smooth isopleths were drawn and values read along  $77\frac{1}{2}^{\circ}\text{E}$  at  $2\frac{1}{2}$  degree latitude intervals from  $5^{\circ}\text{N}$  to  $25^{\circ}\text{N}$ .

At each latitude point, a smooth vertical profile was drawn for  $u$  versus  $p$ . From this profile, the value of  $u$  were picked up at 50 mb interval, from 850 to 150 mb. Thus at each latitude point, we had 15 values along the vertical. We fitted, by the least square method, the parabolic curve,

$$u = a + b \log_{10} p + c (\log_{10} p)^2 \quad \dots \quad (1)$$

where  $u$  is in knots and  $p$  is in mb. At each isobaric level from  $5^\circ\text{N}$  to  $25^\circ\text{N}$ , we had nine values of  $a$ ,  $b$  and  $c$ . To these nine values of  $a$ ,  $b$  and  $c$ , we fitted by the least square method, the curve

$$a = a_0 + a_1\phi + a_2\phi^2 \quad \dots \quad (2A)$$

$$b = b_0 + b_1\phi + b_2\phi^2 \quad \dots \quad (2B)$$

$$c = c_0 + c_1\phi + c_2\phi^2 \quad \dots \quad (2C)$$

where  $\phi$  represents the latitude in degrees.

Thus we had the fitted function :

$$u = (a_0 + a_1\phi + a_2\phi^2) + (b_0 + b_1\phi + b_2\phi^2)(\log_{10} p) + (c_0 + c_1\phi + c_2\phi^2)(\log_{10} p)^2 \quad \dots \quad (3)$$

## 2.2 Goodness of fit

An analysis of variance given in Table 1a for the parabolic curve  $u$  against  $\log_{10} p$  shows that the linear and ~~second~~ degree terms are significant for all latitudes even at 1 percent level.

Similar analysis of variance given in Table 1b for parabolic curve fitted to coefficients  $a$ ,  $b$  and  $c$  against latitude also shows that first



degree and second degree terms are significant for all the coefficients at 1 percent level except for the coefficient  $c$  which is significant at 2.5 percent level.

Table 2 gives observed and fitted zonal winds. It will be seen that the difference between the observed and fitted values rarely exceeds two knots and only at four places it exceeds four knots. Thus, the fitted values agree very well with the observed values. As explained in section 2.3 below the fitting was done by graphical interpolation south of  $5^{\circ}\text{N}$ . All subsequent calculations referred to in this paper for latitude belt  $5^{\circ}\text{N}$  to  $25^{\circ}\text{N}$  were done on the basis of equation 3. The vertical cross section of fitted  $u$  is shown in fig.1. Positive  $u$ 's indicate westerlies and negative  $u$ 's indicate easterlies.

The values of coefficients are given below :

$$\begin{array}{lll} a_0 = -673.20, & a_1 = -56.94, & a_2 = 2.93 \\ b_0 = 426.43, & b_1 = 39.76, & b_2 = -2.03 \\ c_0 = -65.41, & c_1 = -6.73, & c_2 = 0.34 \end{array} \quad (4)$$

### 2.3 Calculation of $u$ south of $5^{\circ}\text{N}$ :

From the calculations of section 2.2 above, we had, at  $5^{\circ}\text{N}$ , the values of  $u$  and  $\frac{\partial u}{\partial y}$  at 50 mb. interval. We also had the observed normal wind values at equator (Gan island, lat.  $0.7^{\circ}\text{S}$ , long.  $73.1^{\circ}\text{E}$ ). We did the graphical interpolation for  $u$  and  $\frac{\partial u}{\partial y}$  between  $5^{\circ}\text{N}$  and the equator, keeping  $\frac{\partial u}{\partial y}$  nearly zero at the equator at levels where meridional component of normal wind is small. Using these interpolated values of  $u$  between equator and  $5^{\circ}\text{N}$  at various isobaric levels, we calculated other parameters using geostrophic relationship through shear wind wherever necessary.



#### 2.4 Calculation of vertical wind shear :

From equation 3, it follows that

$$\frac{\partial u}{\partial p} = \frac{[(b_0 + b_1 \phi + b_2 \phi^2) + 2(c_0 + c_1 \phi + c_2 \phi^2) \log_{10} p] \times \log_{10} e}{p} \text{ ---- (5)}$$

We employed eq.(5) to calculate  $\frac{\partial u}{\partial p}$  between latitudes  $5^\circ \text{N}$  and  $25^\circ \text{N}$ .

Vertical wind shears south of  $5^\circ \text{N}$  were calculated graphically. The vertical wind shear expressed in knots per 100 mb. is shown in Table 3.

#### 2.5 Calculation of temperature :

Here we introduced geostrophic approximation through thermal wind relationship

$$\frac{\partial u}{\partial p} = \frac{R}{f p} \frac{\partial T}{\partial y} \text{ ---- (6)}$$

Hence

$$\frac{\partial T}{\partial y} = \frac{f}{R} [(b_0 + b_1 \phi + b_2 \phi^2) + 2(c_0 + c_1 \phi + c_2 \phi^2) \log_{10} p] \log_{10} e \text{ ---- (7)}$$

Integrating eq.(7) with respect to  $\phi$ , we get

$$T_{\psi, p} - T_{\psi_0, p} = \log_{10} e \cdot r \cdot 2 \pi \left[ -b_0 \cos \psi + b_1 (\sin \psi - \psi \cos \psi) + \right. \\ \left. + b_2 (2 \psi \sin \psi - (\psi^2 - 2) \cos \psi) + 2 \log_{10} p \left\{ -c_0 \cos \psi + \right. \right. \\ \left. \left. + c_1 (\sin \psi - \psi \cos \psi) + c_2 (2 \psi \sin \psi - (\psi^2 - 2) \cos \psi) \right\} \right]_{\psi_0}^{\psi} \text{ (8)}$$

where  $\psi$  is the latitude in radians,  $\phi$  is latitude in degrees and  $\omega$  is the angular velocity of earth and  $r$  is radius of the earth. We took  $\psi_0$  corresponding to latitude  $5^\circ \text{N}$ . The R.H.S. of eq.(8) then gives the latitudinal variation of temperature from  $5^\circ \text{N}$  to any latitude for which eq.(8) is valid. These profiles for various isobaric levels are shown in fig.2. In this diagram,  $5^\circ \text{N}$  has been taken as the reference latitude.

If we know the absolute value of temperature at  $5^\circ \text{N}$  at a pressure



level, we can get through eq.(8) or through fig.2 the value of temperature at any latitude at the same pressure level. The values of temperature at  $5^{\circ}$  N were obtained from analysis of radiosonde observations in that region. From this, we obtained temperature values at various latitudes and at different pressure levels. These were compared with radiosonde temperature values and were found to agree very well. This gives an indirect confirmation that thermal wind relationship which is based on geostrophic relationship, holds very well over the Indian region  $5^{\circ}$  N to  $25^{\circ}$  N for climatological zonal winds in the monsoon season.

Same eq.(6) was used to evaluate temperature at isobaric levels south of  $5^{\circ}$  N. For use of this equation, values of  $\frac{\partial u}{\partial p}$  as determined graphically in section 2.4 were used. The temperature profiles thus derived for latitude south of  $5^{\circ}$  N are also shown in fig.2. Due to extreme closeness of the lines, the graphs are shown only for pressure levels 800, 500 and 150 mb.

## 2.6 Calculation of static stability :

There are various measures of static stability. Here, we present only one measure of static stability given by

$$S = - \frac{\alpha}{\theta} \frac{\partial \theta}{\partial p} = \frac{R}{p^2} \left( K T - \frac{\partial T}{\partial \log_e p} \right) \text{-----} (9)$$

in the usual notation.

This parameter occurs in thermodynamic energy equation and is frequently used in dynamical meteorology. The values of  $T$  were available from section 2.5. The plot of  $T$  against  $\log_e p$  at all latitudes showed a linear relationship between 850 mb. and 150 mb levels of the form :

$$S \approx \frac{R}{p^2} \left( K T - \frac{T_{850} - T_{150}}{\log_e 850 - \log_e 150} \right) \text{-----} (10)$$



The values of  $S$  so obtained are given in Table 4.

#### 2.7 Calculation of $-\frac{\partial u}{\partial y}$ on isobaric surfaces :

From equation (3), we immediately obtained  $-\frac{\partial u}{\partial y}$  on the isobaric surfaces. This is the contribution of zonal current to relative vorticity. Figures (3a) and (3b) give vertical variation of  $-\frac{\partial u}{\partial y}$  and  $f - \frac{\partial u}{\partial y}$  respectively. It is interesting to find from figure 3a that  $-\frac{\partial u}{\partial y}$  is nearly zero at all levels around latitude  $10^\circ \text{N}$ . Further, at levels below 300 mb, this quantity is negative (anticyclonic vorticity) south of  $10^\circ \text{N}$  and positive (cyclonic vorticity) to the north. The opposite holds above 300 mb level.

#### 2.8 Calculation of $-\frac{\partial u}{\partial y}$ on Isentropic surfaces :

For inertial stability of zonal current, we have the criterion  $f - \left(\frac{\partial u}{\partial y}\right)_\theta \geq 0$ , where subscript  $\theta$  denotes differentiation along an isentropic surface (Kuo, 1956). This criterion of Kuo is based on linearised theory of infinitesimal perturbation on a zonal current of finite relative vorticity. The authors are not aware of theoretical extension of this criterion, but it appears plausible to infer from Kuo's work that in his criterion, vorticity arising out of both zonal and meridional motion can perhaps replace the vorticity of the zonal current. In other words, a fluid layer is inertially stable if  $f + \left(\frac{\partial v}{\partial x} - \frac{\partial u}{\partial y}\right)_\theta \geq 0$ . Analysis of normal winds shows that between the equator and latitude  $5^\circ \text{N}$  there is appreciable meridional component also along with the zonal component. Hence  $\frac{\partial v}{\partial x}$  would also contribute significantly to relative vorticity in this region. Elsewhere, i.e. north of  $5^\circ \text{N}$ , the contribution  $\frac{\partial v}{\partial x}$  appears to be small compared to that of  $-\frac{\partial u}{\partial y}$ . Hence it appears reasonable to conclude from Figure 4 that absolute vorticity on an isentropic surface north of  $5^\circ \text{N}$  is positive and hence to conclude that normal flow in the monsoon season is inertially stable, at least

north of latitude  $5^{\circ}$  N.

## 2.9 Calculation of Jet maximum

On inspection of wind between equator and  $25^{\circ}$  N, it was found that the easterly wind maximum lay north of  $5^{\circ}$  N. Hence we could use equation (3) for the analysis of wind maximum. From the condition  $\frac{\partial u}{\partial \phi} = 0$ , one gets the latitude of maximum wind at constant pressure surface as

$$\phi = - \frac{a_1 + b_1 \log_{10} P + c_1 (\log_{10} P)^2}{2 \{ a_2 + b_2 \log_{10} P + c_2 (\log_{10} P)^2 \}} \text{ ----- (11)}$$

Values of this  $\phi$  and the corresponding values of maximum wind are given below.

P (mb)	latitude of maximum wind ( $^{\circ}$ N)	values of easterly max. wind (Kts.)
150	8.80	64.9
200	8.04	42.5
250	5.53	27.5

## 3. Discussion and Conclusions :

3.1 As already stated, there is close agreement between the observed and the fitted zonal winds as well as the temperatures. This suggests two things :

- i) Method of smoothing adopted in this paper is reasonably sound.
- ii) Assumption of geostrophic balance of climatological zonal winds in the region of Indian monsoon appears to be justified.

The method has yielded a detailed structure of Indian monsoon zonal current in a quantitative manner which would be hard to realise without adopting some



smoothing procedure of the type adopted here. With the availability of this detailed quantitative structure, one can study numerous dynamical characteristics of the zonal current of the Indian monsoon. A few of the characteristics are presented below. Further study is in progress.

3.2. Zonal current is inertially stable (section 2.9):

3.3. As further stated in section 2.8, there is appreciable meridional component in the wind south of  $5^\circ \text{N}$ . Hence  $\frac{\partial u}{\partial y}$  does not represent the whole of relative vorticity in this region. However, north of  $5^\circ \text{N}$ , total motion is nearly zonal and hence  $-\frac{\partial u}{\partial y}$  can be taken approximately as the relative vorticity of the mean flow. If the profile of  $(f - \frac{\partial u}{\partial y})$  along  $y$  direction shows a maximum then the zonal current would be barotropically unstable. (Kuo, 1949). Equation (3) assumes  $u$  as a 2nd degree function of  $y$ . In other words,  $\frac{\partial^2 u}{\partial y^2}$  is constant at any given constant pressure level. The values of  $-\frac{\partial^2 u}{\partial y^2}$  in units of  $10^{-11} \text{ m}^{-1} \text{ sec}^{-1}$  at various levels from the region north of  $5^\circ \text{N}$  are given below.

Level (mb)	850	800	700	600	500	400	300	200	150
$-\frac{\partial^2 u}{\partial y^2}$	0.76	0.75	0.71	0.65	0.54	0.36	0.04	-0.56	-1.10

The value of  $\beta$  decreases from  $2.28 \times 10^{-11} \text{ m}^{-1} \text{ sec}^{-1}$  at  $5^\circ \text{N}$  to  $2.07 \times 10^{-11} \text{ m}^{-1} \text{ sec}^{-1}$  at  $25^\circ \text{N}$ . From these values of  $\beta$  and  $-\frac{\partial^2 u}{\partial y^2}$ , it is clear that  $(\beta - \frac{\partial^2 u}{\partial y^2})$  is positive throughout the region. In other words, absolute vorticity monotonically increases with latitude and there is no maximum or minimum in its values in the region under consideration. In other words, the mean zonal current is barotropically stable along  $77\frac{1}{2}^\circ \text{E}$  between  $5^\circ \text{N}$  and  $25^\circ \text{N}$ .



### 3.4 Latitudinal position of easterly jet maximum

We have already shown in section 2.9 that the latitude of the maximum easterly wind at 150 mb level is close to  $9^{\circ}$  N. From the variation in latitude of maximum wind as given by equation (11) in section 2.9 and from the numerical values given in the same section, it is seen that the surface of maximum wind slopes upwards as we go northwards. This is in agreement with the finding of Mokashi (1971) who analysed mean monthly data of individual stations south of  $20^{\circ}$  N.

Inspection of vertical profiles of zonal wind has shown that the easterly wind minimum occurs above 150 mb level but below 100 mb level. In section 2.9, we used equation (11) to locate the position of maximum wind speed at different constant pressure surfaces. Equation (11) is strictly valid at and below 150 mb level. If we extrapolate and apply equation (11) up to 130 mb level, we get maximum easterly wind of 77.2 kts at lat.  $8.97^{\circ}$  N at that level. Combining this with corresponding figures for 200 and 150 mb figures given in section 2.9, one comes to the conclusion that climatologically the maximum easterly wind during July lies somewhere close to  $9^{\circ}$  N.

According to Koteswaram (1958), the summer easterly jet has a maximum around  $15^{\circ}$  N, between 150 and 100 mb levels. At the time of his analysis, the data were sparse and Koteswaram (1958) was the first to identify the existence of easterly jet over the Indian region. The present analysis suggests that the maximum is likely to be found around  $9^{\circ}$  N rather than  $15^{\circ}$  N. But it must be admitted that the horizontal shear in this easterly jet stream is very weak, as seen from figure 1 itself. Hence, the location of the latitude



of maximum wind in the horizontal comes to be more of academic interest at the moment.

### Acknowledgements

The authors have greatly benefited from discussions with their colleagues Mr. C.M.Dixit, Dr. R.N.Keshavamurty and Mr. S.K.Mishra. Mrs. Leela George and Mr. M.B.Gajare gave valuable assistance in computations.

### REFERENCES

- |                |      |   |
|----------------|------|---|
| Charney, J.G.  | 1947 | "Dynamics of long waves in a baroclinic westerly current", Journal of Meteorology, 135-162  |
| Fjortoft, R.   | 1950 | "Application of Integral Theorem in deriving criteria of stability for laminar flows and for baroclinic circular vortex", Geofysiske publikationer, Vol.17, No.6, pp.5-52 |
| Koteswaram, P. | 1958 | "Easterly Jet Stream in the tropics", Tellus, 43-57   |
| Kuo, H.L.      | 1949 | "Dynamic Instability of two dimensional non-divergent flow in a barotropic atmosphere", Journal of Meteorology, 6, 105-122  |
| Kuo, H.L.      | 1956 | "Forced and free meridional circulation in the atmosphere", Journal of Meteorology, 13, 561-568   |
| Mokashi, R.Y.  | 1970 | "Axis of the tropics easterly jet stream over India and Ceylon", Ind.Met.Deptt. Scientific Report No.156  |

Table 1a : Analysis of variance for significance of second degree term (  $u$  versus  $\log_{10} p$  at various latitudes)

Lat.	Source	Sum of squares	Degrees of freedom	Mean square	F-ratio
5° N	Linear term	8635.4	1	8635.4	2828.6
	2nd degree term	273.8	1	273.8	89.7
	Residual	30.6	12	3.0	
	Total	8939.8	14		
10° N	Linear term	9059.9	1	9059.9	3986.2
	2nd degree term	346.5	1	346.5	152.5
	Residual	27.3	12	2.3	
	Total	9433.7	14		
15° N	Linear term	7732.1	1	7732.1	1655.5
	2nd degree term	327.8	1	327.8	70.2
	Residual	56.0	12	4.7	
	Total	8115.9	14		
20° N	Linear term	5412.4	1	5412.4	3528.6
	2nd degree term	90.2	1	90.2	58.8
	Residual	18.4	12	1.5	
	Total	5521.0	14		
25° N	Linear term	1377.6	1	1377.6	1276.1
	2nd degree term	32.1	1	32.1	29.7
	Residual	12.9	12	1.08	
	Total	1422.6	14		



Table 1b : Analysis of variance for significance of second degree term ( a, b and c versus latitude )

	Source	Sum of squares	Degrees of freedom	Mean square	F-Ratio
For coefficient a	Linear term	357547.1	1	357547.1	93.7
	2nd degree term	103088.1	1	103088.1	27.0
	Residual	22895.7	6	3816.0	
	Total	483530.9	8		
for coefficient b	Linear term	165888.2	1	165888.2	60.5
	2nd degree term	49400.0	1	49400.0	18.0
	Residual	16439.9	6	2740.0	
	Total	231728.1	8		
For coefficient c	Linear term	4547.0	1	4547.0	39.0
	2nd degree term	1392.8	1	1392.8	12.0
	Residual	699.2	6	116.5	
	Total	6639.0	8		

Table 2 : Observed and fitted zonal wind component in kts.

	0°	2.5° N	5° N	7.5° N	10° N	12.5° N	15° N	17.5° N	20° N	22.5° N	25° N
	obs. fitted: obs.										



Table 3 : Vertical wind shear in kts per 100 mb

Lat.(deg)/P (mb)	850	800	700	600	500	400	300	200	150
0		-4.0	-1.0	4.9	6.9	10.4	17.1	22.7	22.1
5° N	2.3	2.7	3.7	5.2	7.5	11.2	18.2	34.3	52.3
7.5° N	2.2	2.7	3.7	5.3	7.7	11.6	19.0	35.9	54.9
10° N	2.2	2.6	3.7	5.3	7.7	11.7	19.2	36.4	55.6
12.5° N	2.2	2.6	3.6	5.2	7.5	11.5	18.8	35.5	54.3
15° N	2.1	2.5	3.5	4.9	7.2	10.8	17.7	33.4	51.0
17.5° N	2.0	2.4	3.3	4.6	6.6	9.9	16.0	30.1	45.8
20° N	1.9	2.2	3.0	4.1	5.8	8.6	13.7	25.5	38.6
22.5° N	1.8	2.0	2.6	3.5	4.8	6.9	10.8	19.6	29.4
25° N	1.6	1.8	2.2	2.7	3.6	4.9	7.3	12.6	18.2



Table 4 : Static stability ( $S = - \frac{\alpha}{\theta} \frac{\partial \theta}{\partial p}$ )

( Unit :  $10^{-4} \text{ m}^2 \text{ sec}^{-2} \text{ mb}^{-2}$  )

Lat. (deg)/P (mb)	5° N	10° N	15° N	20° N	25° N
850	139	142	147	153	157
700	191	196	204	213	219
500	324	334	350	370	383
300	667	698	749	808	849
200	1044	1121	1244	1387	1487
150	1411	1554	1785	2052	2236






FIG.1 VERTICAL CROSS-SECTION OF ZONAL WIND (Fitted values)  
IN Kts. +ve VALUES INDICATE WESTERLIES






FIG.2 LATITUDINAL VARIATION OF TEMPERATURE AT DIFFERENT ISOBARIC LEVELS.




FIG. 3 (a) VALUES OF  $-\frac{\partial u}{\partial y}$  ON ISOBARIC SURFACES (UNITS:  $10^6 \text{ Sec}^{-1}$ )






FIG.3(b) VERTICAL CROSS-SECTION OF  $(f - \frac{\partial u}{\partial y})$  ON ISOBARIC SURFACES (UNITS:  $10^{-4} \text{ Sec}^{-1}$ )




FIG. 4 VERTICAL CROSS-SECTION OF  $(f - \frac{\partial u}{\partial y})$  ON ISENTROPIC SURFACES  
(UNITS:  $10^4 \text{ Sec}^{-1}$ )