

RESEARCH REPORT

RR-020




32.181

ANNUAL AND SEMI-ANNUAL THICKNESS OSCILLATIONS
IN THE NORTHERN HEMISPHERE

by

G. C. Asnani & R. K. Verma



INDIAN INSTITUTE OF TROPICAL METEOROLOGY

Ramdurg House
Ganeshkhind Road
Poona - 5, India

January 1975

381181

Annual and Semi-annual thickness oscillations

RR-20 in the Northern-Hemisphere

G.C. Asnani and R.K. Verma



Abstract

1000-850 mb and 1000-300 mb monthly thicknesses have been subjected to harmonic analysis at about 200 stations in the northern hemisphere. Salient features of the annual mean, the annual oscillation and the semi-annual oscillation are presented. In the annual mean state, the warmest temperatures in the eastern hemisphere are found to occur not at the equator but somewhat to the north, around latitude 10°N . It is shown that the heat low which appears over southeast Asia during the summer monsoon is partly due to the permanent warm temperature over this region. Latitude for latitude, the annual mean temperatures do not exhibit much contrast between land and sea, but the land-sea contrast is seen very prominently in the annual oscillation of temperature. In the annual oscillation, largest amplitudes occur at about the same place and at about the same time in the lower troposphere and also in the entire troposphere, which fact suggests that perhaps one and the same physical mechanism is responsible for the diabatic heating in the lower as well as in the middle and upper troposphere, so far as the annual heating cycle on the global scale is concerned. On this scale, a unified parameterisation of diabatic heating through sensible heat transfer at the bottom of the atmosphere and through latent heat release in lower and middle troposphere seems plausible.

It is shown that the annual oscillation of tropospheric temperature over southeast Asia is not spectacularly different from that over the rest of the northern hemisphere in the same latitudes. Question then arises as to what causes the spectacular annual monsoon cycle over southeast Asia. It is suggested that the special features of the annual monsoon of southeast Asia are due to special permanent features of pressure, temperature and wind field over the area rather than due to very special features of the annual temperature oscillation over the area.

1. Introduction

A number of authors have recently analysed the annual wave in the stratosphere and mesosphere (Reed 1962, 1966, Angell and Korshover 1970, Vanloon and Jenne 1970 b). Ebdon (1971) analysed Canton Island data of wind and temperature to determine the relative importance of the quasi-biennial, the annual and the semi-annual oscillations in the equatorial stratosphere and found significant annual oscillation in temperature.

The study of semi-annual oscillation in the stratosphere has been reported by Reed (1964, 1966), Van Loon, Libitzke and Jenne (1972). So far as the troposphere is concerned, Van Loon and Jenne (1969) studied the semi-annual oscillation in respect of zonal wind and temperature in the southern hemisphere. They (Van Loon and Jenne, 1970 a) also extended their study to the Northern-Hemisphere between longitudes 60°E and 120°E. Van Loon has summarised these in good detail for the southern-hemisphere, in Met. Monograph, Vol.13, 1972. A similar diagnostic study for the troposphere of the whole northern-hemisphere was lacking in literature, till Asnani and Misra (1973) and Asnani and Verma (1973) brought out the analysis of the annual and the semi-annual pressure oscillations for the northern-hemisphere, from sea level to 100 mb level.

The thickness of a layer between two constant pressure surfaces is a measure of the mean virtual temperature of the layer. For some purposes, it can be considered as a more useful parameter for studying the temperature distribution in the atmosphere than the actual temperature at constant height or constant pressure levels.

The authors have chosen thicknesses of layers 1000 mb to 850 mb and 1000 mb to 300 mb, primarily to represent the lowest tropospheric layer and also the entire tropospheric layer respectively.

The purpose of the present study is to bring out the salient features of the tropospheric annual and semi-annual temperature oscillations for the whole northern-hemisphere, and to relate these features to the spectacular annual monsoon cycle of southeast Asia.

2. Data and Analysis Procedure

About 200 stations between equator and 80°N were selected. Mean sea level pressure, surface temperature and the geo-potential heights of 850 mb and 300 mb pressure levels at these stations were obtained from the following data sources :

- i) 'Climatological Normals (CLINO) for climat and climat ship stations for the period 1931-1960'. WMO/OMM-No.117. TP.52, 1971 for m.s.l. pressure and surface temperature.
- ii) 'Short period averages for 1951-1960 and provisional average values for climat temp and climat temp ship stations'. WMO/OMM-No.170.TP.82, 1965 for geopotential heights of 850 mb and 300 mb pressure levels.
- iii) Five-year (1967-71) averages were prepared in Theoretical Studies Division of this Institute for 18 additional near-equatorial stations from the monthly values given in 'Monthly Climatic data for the World'. Vols.20-24, published by WMO in co-operation with U.S. Weather Bureau.
- iv) 'Normals of climat temp based on morning and afternoon/evening radiosonde data for the period 1951-1970', India Meteorological Department, 1972, for 13 Indian stations. This publication gives surface pressure at the station. This station level pressure was first reduced to mean sea level pressure, assuming constant lapse rate of $6.5^{\circ}\text{C. Km}^{-1}$ between the station level and the sea level.

In the first stage, at each station, the monthly mean values of sea level pressure were converted into the monthly mean

values of 1000 mb height, using the relationship,

$$Z_{1000 \text{ mb}} = \frac{RT}{g} \ln \frac{P_{\text{sea level}}}{1000 \text{ mb}}$$

where,

R = Specific Gas constant.

T = Mean virtual temperature of air between sea level and 1000 mb.

g = Acceleration of gravity.

As an approximation, T was taken to be the station level temperature. Analysis showed that the error introduced by this approximation would not affect our results to any significant extent. Station level temperature was readily available in the publications which served as our data sources. Monthly mean values of 1000-850 mb thickness and 1000-300 mb thickness were then obtained at each station. These monthly mean thicknesses were subjected to harmonic analysis to get annual mean as also the amplitude and phases of 12-monthly and 6-monthly oscillations at individual stations. These parameters were plotted on northern hemispheric charts and hand-analysed. The analyses of the annual mean, the 12-monthly oscillation and the 6-monthly oscillation are presented. Since the data consist of normals commencing from January and ending in December, the zero of the phase angle was counted from mid January.

3. Results and Discussion

3.1 Annual Mean State

(a) 1000-850 mb thickness - Annual mean thickness (gpm) of the layer between 1000 mb and 850 mb is shown in Fig. 1(a). Broadly speaking, the pattern is characterised by lower thickness values near the pole (~ 1250 gpm) increasing towards south (> 1400 gpm near the equator). However, one interesting feature deserves special mention. In the eastern hemisphere, the maximum values of thickness i.e. warmest temperatures occur not at the equator but somewhat to the north of the equator.

Fig.1(a), Fig.1(b) and Fig.1(c)

Curiously enough, the warmest region in this annual mean chart has almost the same alignment as the summer monsoon trough in the lower troposphere. For comparison, we show in Fig. 1(b), the position of the summer monsoon trough at 1000, 850, 700 and 500 mb levels during the month of July and also the position of the maximum thickness line for the layer 1000-850 mb (annual mean). It is clear that the heat 'low' which appears over the Indo-Gangetic plain in the lower troposphere during the summer monsoon season is partly due to the permanent warm temperatures over this region in the lower troposphere.

(b) 1000-300 mb thickness - Fig. 2 depicts this pattern. As in case of 1000-850 mb thickness, lower thickness values (8600 gpm) occur near the pole, increasing towards the south (9600 gpm near the equator). Again, the noteworthy feature is observed i.e. in the eastern hemisphere, the maximum values of thickness (or the warmest temperatures) occur not at the equator but somewhat north of the equator.

Fig. 2

3.2 Annual-Oscillation

(a) 1000-850 mb thickness - Fig. 3(a) shows the amplitude of 12-monthly oscillation in the thickness of 1000-850 mb layer. These amplitudes are lowest (< 10 gpm) near the equator and increase northwards upto about 60°N . There are two distinct maxima around 60°N latitude circle, one near 120°E (NE Asia, 100 gpm) and the other near 120°W (NW Canada, 80 gpm). Fig.3(b), shows the time of occurrence of maximum thickness. It is seen that the largest values of the oscillation seen over NE Asia as well as over NW Canada occur during later part of July.

Fig.3(a) and Fig. 3(b)

(b) 1000-300 mb thickness - Fig. 4(a) shows the amplitude of 12-monthly oscillation in the thickness of 1000-300 mb layer. Here also, as in Fig.3(a), the amplitude of the oscillation is smallest (< 50 gpm) near the equator and increases

northwards upto about 60 N. Around this latitude circle, there are two maxima, one over NE Asia (500 gpm) and the other over NW Canada (350 gpm). The peak values of this oscillation also occur during later part of July. The phase analysis is shown in Fig. 4(b).

Fig.4(a) and Fig.4(b)

Figures 3 and 4 bring out the following features :

i) Along a latitude circle, amplitude of the annual oscillation is smaller over the oceanic regions (Atlantic and Pacific) and larger over the continents (Eurasia and North-America). Between the land masses, the oscillation is larger over Eurasian land mass. As a consequence, in the mid-summer, the troposphere is warmer over land masses than over oceanic regions. In mid-winter, the reverse takes place. It is somewhat surprising that this land-sea contrast is not seen so much in the annual mean thickness charts of the lower troposphere (Fig.1(a)) and of the entire troposphere (Fig.2) but is very well marked in the 12-monthly oscillation of the lower troposphere as well as the entire troposphere. Obviously, the lower, middle and the upper troposphere, all get more heated during the summer and more cooled during the winter over the continental land mass than over the oceanic region.

ii) It is generally believed that sensible heat transfer at the earth-air interface is largely responsible for temperature oscillations in the lower troposphere and that the release of latent heat during condensation of water vapour is responsible for heating in the middle and upper troposphere. Figs.3(b) and 4(b) indicate the time when the maximum in the annual oscillation is reached in respect of 1000-850 mb and 1000-300 mb thicknesses respectively. It is seen from these figures, that in the lower troposphere, maximum thickness occurs practically over the same region and almost at the same time. We propose to undertake a more detailed study of the thickness oscillation for smaller layers, like 850-700 mb, 700-500 mb, 500-300 mb etc. to see the

spatial and temporal progression of thickness maxima for various layers. Their space and time relationships will indicate how strong is the vertical coupling between the various layers. If Fig.s 3 and 4 are any indication, it would appear that there is perhaps one and the same physical mechanism responsible for the annual temperature oscillations at various levels. If the physical processes like diffusion of sensible heat from below and release of latent heat in the middle troposphere be really different and unconnected, then there is no reason to anticipate such similarity as we find in Fig's. 3 and 4, unless the apparently different processes are in fact mutually linked very intimately. If so, we may attempt a unified parameterization of diabatic heating through sensible heat transfer and latent heat release, so far as annual cycle on the global scale is concerned. The other possibility is that out of the two physical processes, sensible heat transfer and latent heat release, only one process is dominant for this annual oscillation.

iii) What causes the spectacular annual monsoon cycle of southeast Asia as distinct from the rest of the world ? This question is discussed below with reference to the computational results presented above.

a) Thickness oscillation is another representation of temperature oscillation. Temperature oscillation arises from horizontal temperature advection, diabatic heating and vertical motion. Of these, diabatic heating appears to be playing a dominant role. Whatever be the relative role of these mechanisms of heating, figure 3(a) clearly show that in the lower troposphere (1000-850 mb layer), air over southeast Asia gets heated during summer about as much as the air elsewhere at the same latitude, say over north America. In Fig.3(a), isopleths of amplitude 10 to 60 gpm are about similarly placed over southeast Asia as over north America between latitudes 10°N and 40°N . Hence, the lower tropospheric summer 'heat low' of south Asia which induces the air to 'flow into' the land area of south Asia during summer and causes the summer monsoon as distinct from the summer monsoon

elsewhere could not be due to any excessive summer heating over southeast Asia. The fact is that the annual mean patterns of lower tropospheric temperature (Fig.1 a) and pressure (Fig.1 c, taken from Asnani and Mishra, 1973) over southeast Asia are somewhat different from these over the rest of the world. For example, there is a permanent warm lower troposphere over southeast Asia as shown in Fig. 1(a) and discussed in section 3.1(a) above. There is also a permanent low pressure trough over north India as seen in Fig. 1(c). The ultimate causes of these special permanent features of the lower tropospheric temperature and pressure patterns over southeast Asia obviously are the special orography and land-sea contrasts in the region. It is not the purpose of this paper to go into a discussion of these ultimate causes of permanent features. What is aimed to be brought out here is that the annual oscillation of lower tropospheric temperature over southeast Asia is not spectacularly different from that over the rest of the world in the same latitudes.

b) 1000-300 mb thickness oscillation shown in Fig.4(a) roughly represents the temperature oscillation of the whole troposphere. Here, the maximum amplitude of 500 gpm occurs over northeast China and neighbourhood, outside the region of southeast Asian monsoon. It will be seen that south of 30°N , the amplitudes of the annual oscillation over southeast Asia are not very different from these over the rest of the world in the same latitudes. The amplitudes are quite comparable, latitude for latitude. Here also, one cannot say that the whole troposphere over southeast gets much more warmed up than elsewhere and hence the spectacular summer monsoon of southeast Asia. In fact, the summer heating of the troposphere in southeast Asia is about the same as over the rest of the world in the same latitudes.

c) What appears to be special about southeast Asia is that due to special orographic features of this region, the permanent tropospheric pressure, temperature and flow patterns have some special peculiarities. Annual oscillation of tropospheric temperature is not very different over southeast Asia in

comparison to the rest of the world, latitude for latitude. When this annual oscillation is superimposed on the permanent annual mean patterns, the resultant flow pattern in southeast Asia has some special features like deep westerly current lying under a broad easterly jet stream in the upper troposphere, etc.

Land-sea contrast and orography in SE Asia is also such that the resultant lower tropospheric winds in summer pick up considerable moisture over the sea surface and dump it over land areas in the form of extensive cloud and rain. This phenomenon of cloud and rain in summer occurs over other parts of the world also but with much less intensity than over southeast Asia.

iv) The maximum amplitude of oscillation (Fig.4 a) is larger (500 gpm) over north Asia than over north America (360 gpm). Apparently, the maximum amplitude of oscillation depends upon the horizontal size of the land mass. Eurasian land mass is larger than north American land mass. The dependency of the amplitude of oscillation on the horizontal extent of the source of heating and cooling was first pointed out by Jeffreys (1926) and has been recently emphasised by Jecques and Wiin-Nielsen (1971) and Asnani and Mishra (1973).

3.3 Semi-Annual Oscillation

Fig's. 5 and 6(a) show the amplitude of semi-annual oscillation in the thicknesses of 1000-850 mb layer and 1000-300 mb layer respectively. In the case of 1000-850 mb thickness the amplitudes are smallest (< 2 gpm) near the equator and largest (> 10 gpm) in the polar latitudes. Phases are not presented as they were found to be highly variable, presumably due to very small amplitudes.

Fig.5, Fig.6(a) and Fig.6(b)

In the case of 1000-300 mb thickness, the lowest amplitudes are near the equator (less than 10 gpm) and largest amplitudes are in the polar latitudes (greater than 60 gpm and even greater than 100 gpm in some regions). Within this general distribution, two secondary maxima are observed in the sub-tropical latitudes, one over U.S.A. and the other over the Mediterranean. The semi-annual oscillations in tropical and higher latitudes are in opposite phase. In higher latitudes maxima occur during January and July, while in tropics maxima occur during April and October.

4. Conclusions

(i) In the annual mean, the warmest temperatures in the eastern hemisphere are found to occur not at the equator but somewhat to the north along latitude 10 N. It is shown that the heat low which appears over southeast Asia during the summer monsoon is partly due to the permanent warm temperatures over this region.

(ii) Latitude for latitude, the annual mean temperatures do not exhibit much contrast between land and sea, but the land-sea contrast is seen very prominently in the annual oscillation of temperature.

(iii) In the annual temperature oscillation, largest amplitudes occur at about the same place and at about the same time in the lower troposphere and also in the entire troposphere. This suggests that perhaps one and the same physical mechanism is responsible for the diabatic heating in the lower as well as in the middle and upper troposphere, so far as the annual heating cycle, on the global scale, is concerned. On this scale, unified parameterisation of diabatic heating through sensible heat transfer at the bottom of the atmosphere and through latent heat release in lower and middle troposphere would appear plausible.

(iv) It is shown that the annual oscillation of tropospheric temperature over southeast Asia is not spectacularly different from that over the rest of the northern hemisphere in the same latitudes. Question then arises as to what causes the spectacular annual monsoon cycle over southeast Asia. It is suggested that the special features of the annual monsoon of southeast Asia are due to special permanent features of pressure, temperature and wind field over the area rather than due to very special features of the annual temperature oscillation over the area.

Acknowledgements

Computational and plotting help was given by Miss Khade and Mrs. George. Diagrams were drawn by Mr. Rahalkar and Mr. Gade. Typing was done by Mr. Goklaney. Their help is gratefully acknowledged.

REFERENCES

- | | | |
|-----------------------------------|------|---|
| Angell, J.K., and
J. Korshover | 1970 | 'Quasi-binennial, annual and Semi-annual zonal wind and temperature harmonic amplitudes and phases in the stratosphere and low mesosphere of the northern hemisphere', J.Geophys.Res., 75, 543-550. |
| Asnani, G.C., and
Mishra, S.K. | 1973 | 'Annual pressure oscillation from sea level to 100 mb in the northern hemisphere', paper presented at NWP symposium, New Delhi, 25,27 July 1973. |
| Asnani, G.C. and
Verma, R.K. | 1973 | 'Semi-annual pressure oscillation from sea level to 100 mb in the northern hemisphere', RR-017, Indian Institute of Tropical Meteorology, Poona. |

- | | | |
|--|--------|---|
| Ebdon, R.A. | 1971 | 'Periodic fluctuations in equatorial stratospheric temperature and wind'. Met.Magazine, 100, 84-90. |
| Jacques, D. and Wiin-Nielson, A. | 1971 | 'Response of a middle-latitude model atmosphere to forcing by topography and stationary heat sources', Mon.Wea.Rev., 99, 564-576. |
| Jeffreys, H. | 1926 | 'On the dynamics of geostrophic winds', Quart.J.Roy.Met.Soc., 52, 85-104. |
| Reed, R.J. | 1962 | 'Some features of the annual temperature regime in the tropical stratosphere', Mon.Weather Rev., 90, 211-215. |
| Reed, R.J. | 1964 | 'A climatology of wind and temperatures in the tropical stratosphere between 100 mb and 10 mb', Contract NWRP 26-0564-092, 56 pp., U.S., Navy Weather Res. Facil., Norfolk, Va. |
| Reed, R.J. | 1966 | 'Zonal wind behaviour in the equatorial stratosphere and lower mesosphere', J.Geophys.Res., 71, 4223-4233. |
| VanLoon, Harry and Jenne, Roy L. | 1969 | 'The half-yearly oscillations in the tropics of the southern hemisphere', J.Atmos.Sci., 26, 218-232. |
| VanLoon, Harry and Jenne, Roy L. | 1970 a | 'On the half-yearly oscillations in the tropics', Tellus, 22, 391-398. |
| VanLoon, Harry and Jenne, Roy L. | 1970 b | 'The annual wave in the temperature of the low stratosphere', J.Atmos. Sci., 27, 701-705. |
| VanLoon, Harry, Labitzke, K. and Jenne, Roy L. | 1972 | 'Half-yearly wave in the stratosphere', J.Geophys.Res., 77, 3846-3855. |
| VanLoon, et al. | 1972 | 'Meteorology of the southern hemisphere', Met. Monographs, 13, Amer. Met. Soc. |

000000

Legend of diagrams

- Fig. 1(a) : Annual mean thickness (gpm) of 1000-850 mb layer. The dash-dot line is the line of largest thickness.
- Fig. 1(b) : Position of monsoon trough at 1000, 850, 700 and 500 mb levels (dotted lines) during July and that of maximum thickness line for the 1000-850 mb layer (continuous line).
- Fig. 1(c) : Annual normal sea level pressure (mb).
- Fig. 2 : Annual mean thickness (gpm) of 1000-300 mb layer. The dash-dot line is the line of largest thickness.
- Fig. 3(a) : Amplitude (in gpm) of 1000-850 mb thickness in annual oscillation.
- Fig. 3(b) : Time of occurrence of maximum thickness (1000-850 mb) in annual oscillation.
- Fig. 4(a) : Amplitude (in gpm) of 1000-300 mb thickness in annual oscillation.
- Fig. 4(b) : Time of occurrence of maximum thickness (1000-300 mb) in annual oscillation.
- Fig. 5 : Amplitude (in gpm) of 1000-850 mb thickness in semi-annual oscillation.
- Fig. 6(a) : Amplitude (in gpm) of 1000-300 mb thickness in semi-annual oscillation.
- Fig. 6(b) : Time of occurrence of maximum thickness (1000-300 mb) in semi-annual oscillation.

nnnnnnnn




FIG.-1(a): ANNUAL MEAN THICKNESS (gpm) OF 1000-850 mb LAYER. THE DASH-DOT LINE IS THE LINE OF LARGEST THICKNESS.




FIG.-1(b): POSITION OF MONSOON TROUGH AT 1000, 850, 700 AND 500 mb LEVELS (DOTTED LINES) DURING JULY AND THAT OF MAXIMUM THICKNESS LINE FOR THE 1000-850 mb LAYER (CONTINUOUS LINE).







FIG.-1(c): ANNUAL NORMAL SEA LEVEL PRESSURE (mb)






FIG.-4(a): AMPLITUDE (IN gpm) OF 1000-300 mb THICKNESS IN ANNUAL OSCILLATION.




FIG.- 4(b): TIME OF OCCURRENCE OF MAXIMUM THICKNESS(1000-300mb) IN ANNUAL OSCILLATION.




FIG.- 5 : AMPLITUDE (IN gpm) OF 1000-850 mb THICKNESS IN SEMI-ANNUAL OSCILLATION.




FIG.-6(a): AMPLITUDE (IN gpm) OF 1000-300 mb THICKNESS IN SEMI-ANNUAL OSCILLATION.




FIG.-6(b): TIME OF OCCURRENCE OF MAXIMUM THICKNESS (1000-300mb) IN SEMI-ANNUAL OSCILLATION.